



## Temperature Management in High Tunnels

*Brent Black*, Extension Fruit Specialist, and *Dan Drost*, Extension Vegetable Specialist

### Introduction

High tunnels have proven to be an effective method for extending the growing season for a number of high-value crops in a diverse range of climates. High tunnels are large plastic covered structures that are used to modify the growing conditions of the covered area. High tunnels are similar to greenhouses, except that warming and cooling of high tunnel is usually entirely passive, where expensive cooling fans and heaters are often used in greenhouses.

High tunnels are used for cold protection, and to provide optimal growing conditions for longer periods of time. The key to tunnel management is to understand the temperature requirements of the crop. This fact sheet spells out temperature considerations and how these are best addressed using high tunnels and other season-extending technologies.




Figure 1. Typical relationship between temperature and plant growth rate.

### Temperatures and Growth

Figure 1 shows the typical relationship between temperature and plant growth. At the left side of the graph is a point where temperatures are cold enough to

cause plant injury. When temperatures are above that point, the plant may survive un-injured, but may not grow until temperatures exceed the minimum or base temperature for growth. As temperatures increase above the base, the plant growth rate also increases up to the point where temperature is optimal and plant growth is maximized. As temperatures exceed the optimum, then plant growth slows until the point where heat injury begins to take place.

The “critical” or “cardinal” temperatures are the minimum, optimum and maximum temperatures that define this growth curve, and these temperatures differ from crop to crop. A cool season crop like lettuce can withstand colder temperatures than a warm season plant like watermelon. Cool season plants also have a lower optimum temperature than do warm season crops. Therefore, they tend to undergo heat stress symptoms at lower temperatures. The cardinal temperature for seed germination and plant growth of some common fruit and vegetable crops is shown in Tables 1-3.

**Table 1. Cardinal temperature for fruit crop growth.**

| Crop       | Minimum<br>(°F) | Optimum<br>(°F) | Maximum<br>(°F) |
|------------|-----------------|-----------------|-----------------|
| Strawberry | 40              | 65-75           | 85              |
| Raspberry  | 40              | 65-75           | 90              |
| Blackberry | 40              | 75-85           | 95              |

### Temperatures Management

High tunnels are heated when sunlight warms the plants and soil. The warmed plants and soil reradiate the heat, warming the air within the tunnel. Temperatures within a closed tunnel can increase rapidly on sunny days even when outside air temperatures are relatively cold. Even in late winter, it is often necessary to ventilate tunnels on sunny days to prevent the temperatures from exceeding the crop’s growth optimum. One of the key management tools is to situate

**Table 2. Cardinal temperature for seed germination of select vegetables.**

| Crop         | Minimum (°F) | Optimum (°F) | Maximum (°F) | Ideal range (°F) |
|--------------|--------------|--------------|--------------|------------------|
| Asparagus    | 50           | 75           | 95           | 60-85            |
| Bean         | 60           | 80           | 95           | 60-85            |
| Bean, lima   | 40           | 85           | 85           | 65-85            |
| Beet         | 40           | 85           | 95           | 50-85            |
| Cabbage      | 40           | 85           | 100          | 45-95            |
| Carrot       | 40           | 80           | 95           | 45-85            |
| Cauliflower  | 40           | 80           | 100          | 45-85            |
| Celery       | 40           | 70           | 85           | 60-70            |
| Chard, Swiss | 40           | 85           | 95           | 50-85            |
| Corn         | 50           | 95           | 105          | 60-95            |
| Cucumber     | 60           | 95           | 105          | 60-95            |
| Eggplant     | 60           | 85           | 95           | 75-90            |
| Lettuce      | 35           | 75           | 85           | 40-80            |
| Muskmelon    | 60           | 90           | 100          | 75-95            |
| Okra         | 60           | 95           | 105          | 70-95            |
| Onion        | 35           | 75           | 95           | 50-95            |
| Parsley      | 40           | 75           | 90           | 50-85            |
| Parsnip      | 35           | 65           | 85           | 50-70            |
| Pea          | 40           | 75           | 85           | 40-75            |
| Pepper       | 60           | 85           | 95           | 65-95            |
| Pumpkin      | 60           | 90           | 100          | 70-90            |
| Radish       | 40           | 85           | 95           | 45-90            |
| Spinach      | 35           | 70           | 85           | 45-75            |
| Squash       | 60           | 95           | 100          | 70-95            |
| Tomato       | 50           | 85           | 95           | 60-85            |
| Turnip       | 40           | 85           | 105          | 60-105           |
| Watermelon   | 60           | 95           | 105          | 70-95            |

From Knott's Handbook for Vegetable Growers (4<sup>th</sup> Edition).

sufficiently reduce the temperature then the side walls can also be opened (Figure 2). Doors should be closed in the late afternoon to retain heat through the night time. Low tunnels placed over individual rows of crops within the high tunnel can add an additional “temperature lift” to prevent cold damage at night (Figure 3). Low tunnels also require careful ventilation to prevent extreme high temperatures. Without proper ventilation of both high and low tunnels, temperatures in a Northern Utah high tunnel have exceeded 120°F on a sunny day in early March.



**Figure 3. Low tunnels within a high tunnel.**

Later in the spring, as the day and night temperatures remain above the minimum for the crop, the plastic can be removed from the tunnel and replaced with shade cloth to reduce the risk of sunburn or other heat stress.

**Table 3. Cardinal temperature for growth of selected vegetables.**

| Crop                                   | Min. (°F) | Opt. (°F) | Max. (°F) |
|--|-----------|-----------|-----------|
| Chive, Garlic, Leek, Onion             | 45        | 55-75     | 85        |
| Beet, Broad Bean, Broccoli, Cabbage    |           |           |           |
| Chard, Collards, Kale, Parsnips        | 40        | 60-65     | 75        |
| Radish, Spinach, Turnip                |           |           |           |
| Artichoke, Carrot, Cauliflower, Celery |           |           |           |
| Endive, Lettuce, Mustard,              | 45        | 60-65     | 75        |
| Parsley, Pea, Potato                   |           |           |           |
| Lima Bean, Snap Bean                   | 50        | 60-70     | 80        |
| Sweet corn                             | 50        | 60-75     | 95        |
| Pumpkin, Squash                        | 50        | 65-75     | 90        |
| Cucumber, Muskmelon                    | 60        | 65-75     | 90        |
| Sweet Pepper, Tomato                   | 65        | 70-75     | 80        |
| Eggplant, Hot Pepper, Okra,            | 65        | 70-85     | 95        |
| Sweet Potato, Watermelon               |           |           |           |

From Knott's Handbook for Vegetable Growers (4<sup>th</sup> Edition).

an accurate minimum-maximum thermometer in the house and use it to track daily low and high temperatures. Carefully following tunnel temperatures is essential to developing management strategies for maintaining optimum plant growth conditions.

Opening the end doors or vents is the first step in ventilation and temperature regulation. If this does not



**Figure 2. Side ventilation of a tomato high tunnel.**

Utah State University is committed to providing an environment free from harassment and other forms of illegal discrimination based on race, color, religion, sex, national origin, age (40 and older), disability, and veteran's status. USU's policy also prohibits discrimination on the basis of sexual orientation in employment and academic related practices and decisions.

Utah State University employees and students cannot, because of race, color, religion, sex, national origin, age, disability, or veteran's status, refuse to hire; discharge; promote; demote; terminate; discriminate in compensation; or discriminate regarding terms, privileges, or conditions of employment, against any person otherwise qualified. Employees and students also cannot discriminate in the classroom, residence halls, or in on/off campus, USU-sponsored events and activities.

This publication is issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Noelle E. Cockett, Vice President for Extension and Agriculture, Utah State University.